Dermal Filler Post-Treatment Instructions

- Swelling, bruising, tenderness, numbness, and areas of firm nodules may occur.
- Cosmetic fillers are long lasting, but not permanent. Longevity depends on the areas treated and your body's metabolism.
- DO NOT APPLY ANYTHING TO THE SKIN UNTIL THE DAY AFTER TREATMENT: No cleanser, moisturizer, or makeup!
- DO NOT massage, touch, or manipulate the injection site. Avoid heavy exercise the day of your treatment.
- Ice packs may be used to the treated area during the first 12 hours. Ice for 15 minutes on every hour.
- Avoid dental work for 30 days after filler injections. Since the potential risk for infection is increased.
- Avoid sleeping on your side or stomach for 24 hours post-treatment to prevent dispersing the product. We recommend you sleep elevated on your back and use pillows to stabilize yourself.
- If experiencing swelling, sleep with head elevated for 2-3 days to decrease swelling.
- You may take acetaminophen/Tylenol if you experience any mild tenderness or discomfort.
- Wait a minimum of four weeks (or as directed by your provider) before receiving any skin care or laser treatments.
- For most patients, the benefits of dermal fillers can be enhanced by using a medical-grade skincare system. Please contact your provider or a member of the staff for product recommendations.
- If you experience bruising, Arnica Montana natural supplement can be taken to help bruising to resolve at a faster rate; take 5 tablets three times per day (this can be started 1 week prior to injections).
- If you experience any of the following symptoms, contact our office immediately: dusky or white discoloration of injected areas, mottling or unusual bruising, severe or increasing pain, redness, increasing warmth or coolness to touch, severe swelling, itching, blisters, difficulty swallowing or smiling, fever, or chills.

If you notice onset of any of these symptoms,	s, call our office immediately at 732-380-6010 to spea
with a provider or our answering service.	

Please Sign acknowledgement of the post care instructions above.

Patient Sign:		